

## Durham Research Online

---

### Deposited in DRO:

12 June 2018

### Version of attached file:

Published Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Carreras, Javier and Livendahl, Madeleine and McGonigal, Paul R. and Echavarren, Antonio M. (2014) 'Gold(I) as an artificial cyclase : short stereodivergent syntheses of (-)-epiglobulol and (-)-4,7- and (-)-4,7-aromadendranediols.', *Angewandte chemie international edition.*, 53 (19). pp. 4896-4899.

### Further information on publisher's website:

<https://doi.org/10.1002/anie.201402044>

### Publisher's copyright statement:

© 2014 The Authors. Published by WileyVCH Verlag GmbH Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

### Additional information:

---

### Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

# Gold(I) as an Artificial Cyclase: Short Stereodivergent Syntheses of (–)-Epiglobulol and (–)-4β,7α- and (–)-4α,7α-Aromadendranediols\*\*

Javier Carreras, Madeleine Livendahl, Paul R. McGonigal, and Antonio M. Echavarren\*

**Abstract:** Three natural aromadendrane sesquiterpenes, (–)-epiglobulol, (–)-4β,7α-aromadendranediol, and (–)-4α,7α-aromadendranediol, have been synthesized in only seven steps in 12, 15, and 17% overall yields, respectively, from (E,E)-farnesol by a stereodivergent gold(I)-catalyzed cascade reaction which forms the tricyclic aromadendrane core in a single step. These are the shortest total syntheses of these natural compounds.

Aromadendranes are a family of hydroazulenes named after (+)-aromadendrene (**1**, Figure 1), the main component in the essential oil from *Eucalyptus* trees. The related sesquiterpenoids (–)-globulol (**2**), (–)-epiglobulol (**3**), (–)-4α,7α-aromadendranediol (**4**), and (–)-4β,7α-aromadendranediol (**5**)




Figure 1. Naturally occurring aromadendranes.

[\*] Dr. J. Carreras, M. Livendahl, Dr. P. R. McGonigal, Prof. A. M. Echavarren  
Institute of Chemical Research of Catalonia (ICIQ)  
Av. Paisos Catalans 16, 43007 Tarragona (Spain)  
Prof. A. M. Echavarren  
Departament de Química Analítica i Química Orgànica  
Universitat Rovira i Virgili, C/Marcel·lí Domingo s/n  
43007 Tarragona (Spain)  
E-mail: aechavarren@iciq.es

[\*\*] We thank the European Research Council (Advanced Grant No. 321066), the MINECO (CTQ2010-16088/BQU), and the ICIQ Foundation for financial support. We thank Elina Buitrago (visiting student from the University Stockholm) for experiments on the synthesis of (±)-**3**, and the ICIQ X-Ray Diffraction and Chromatography units.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201402044>.


© 2014 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

are widespread in plant species<sup>[1]</sup> and display antifungal,<sup>[2]</sup> antibacterial,<sup>[3]</sup> antiviral,<sup>[4]</sup> cytotoxic,<sup>[5]</sup> and other activities.<sup>[6]</sup> Interestingly, the antipodes of **1** and other aromadendrenes have been isolated from corals.<sup>[7]</sup> Aromadendranes with amino, isonitrile, isothiocyano, and urea functionalities at C4 have been found in sponges.<sup>[8]</sup> Diterpenoids with an aromadendrane structure are also natural products.<sup>[9]</sup>

The synthesis of members of this family of tricyclic sesquiterpenes has attracted significant interest.<sup>[10]</sup> (–)-Epiglobulol (**3**), isolated in hop<sup>[11]</sup> and many essential oils,<sup>[12]</sup> was prepared from **1** or the corresponding ketone (apoaromadendrone).<sup>[13]</sup> A first total synthesis of **3** from the chiral pool was accomplished in eight steps (4% overall yield).<sup>[14]</sup> A recent synthesis of (±)-epiglobulol in 18 steps used a rhodium(I)-catalyzed hydroacylation/cycloisomerization as the key step.<sup>[15]</sup>

(–)-4α,7α-Aromadendranediol (**4**) was isolated from a marine coral *Sinularia may*<sup>[7]</sup> and the leaves of the Amazonian tree *Xylopia brasiliensis*.<sup>[2]</sup> A semisynthesis of **4** from (+)-spathulenol<sup>[7]</sup> and one total synthesis have been reported.<sup>[16]</sup> This total synthesis involved a three-reaction sequence in a three-component reaction to generate four stereogenic centers in one step and required ten steps to produce **4** in 23% overall yield. (–)-4β,7α-Aromadendranediol (**5**) has been isolated from the leaves of *Chloranthus glaber*.<sup>[17]</sup> A semisynthesis of **5** from (+)-spathulenol has been reported.<sup>[7]</sup>

We developed a gold(I)-catalyzed cascade cyclization of the diyne **6**, a cascade consisting of a cyclization, 1,5-migration of the propargylic OR group, and intramolecular cyclopropanation, thus leading to tricyclic structures closely related to the aromadendrene sesquiterpenes (Scheme 1).<sup>[18]</sup> This reaction is stereospecific since (E)-**6** gave the tricyclic product **7** having the relative configuration of **3** and **5**, whereas the geometrical isomer of **6** led to **8**, the C4 epimer of **7**, having the configuration of **2** and **4**. We recently applied




Scheme 1. Gold-catalyzed formation of tricyclic cores of the aromadendranes by cyclization/1,5-OR migration/intramolecular cyclopropanation.

a strategy based on a gold(I)-catalyzed cyclization/1,5-OR migration/intermolecular cyclopropanation for the first total synthesis of (+)-schisanwilsonene **A**. As part of our program on the synthesis of terpenoids by using new gold-catalyzed cyclization cascades,<sup>[19]</sup> we decided to target **3**, **4**, and **5**, each of which present six stereogenic centers in a tricyclic skeleton. In principle, **3** and **5** could be synthesized from the dienyne (*S,E*)-**6** (Scheme 1), whereas **4** would be prepared from geometric isomer (*S,Z*)-**6**. However, although enantioenriched (*E*)-**6** could be readily prepared from (*E,E*)-farnesol (**9**), the starting material, (*E,Z*)-farnesol, required for the synthesis of (*Z*)-**6** is not commercially available.<sup>[20]</sup>

Herein we report a simple solution to this problem and it allows general access to this class of sesquiterpenes from (*S,E*)-**6** as a common precursor by means of a stereodivergent gold(I)-catalyzed cascade process. The reaction can take place intramolecularly by 1,5-migration of OR in **A** and in the presence of an external nucleophile (via **B**), thus leading to **7** and **8**, respectively, having opposite configurations at C4 (Scheme 1). Starting from (*R,E*)-**6**, enantiomeric aromadenranes can be similarly obtained. This proposal is based on our initial mechanistic study in the *Z* series, in which we found that the cyclopropyl gold(I) carbene intermediate could be trapped with methanol to form an epimeric compound as a minor product.<sup>[18]</sup> In these transformations, gold(I) acts as an artificial cyclase,<sup>[21]</sup> thus mimicking the action of terpene cyclases forming polycyclic skeletons by the selective activation of the alkyne terminus of a dienyne, to readily build a tricyclic skeleton with exquisite stereocontrol.<sup>[22,23]</sup>


The dienyne (*S,E*)-**6** (*R* = Bn) was prepared in four steps and 62% overall yield by using a route similar to that used in the transformation of the lower homologue geraniol.<sup>[19a,24]</sup> the transformation involved the known Sharpless asymmetric epoxidation of (*E,E*)-farnesol (**9**) to give the epoxide (*S,S*)-**10** (88% yield, 91:9 e.r.)<sup>[25,26]</sup> (Scheme 2). Substitution of the



**Scheme 2.** a) L-(+)-DIPT, Ti(OiPr)<sub>4</sub>, tBuOOH, 4 Å M.S., CH<sub>2</sub>Cl<sub>2</sub>, −48 °C, 88%, 82% ee;<sup>[25]</sup> b) PPh<sub>3</sub>, NaHCO<sub>3</sub>, CCl<sub>4</sub>, reflux, 6 h, 94%; c) *n*BuLi, THF, −40 °C, 2 h, 82%; d) BnBr, NaH, Bu<sub>4</sub>NI, THF, 23 °C, 12 h, 91%. DIPT = diisopropyl tartrate, M.S. = molecular sieves, THF = tetrahydrofuran.

primary alcohol by chloride with CCl<sub>4</sub> and PPh<sub>3</sub> gave **11**, which was treated with *n*BuLi to yield the propargylic alcohol **12**. Finally, benzylation under standard reaction conditions gave (*S,E*)-**6**.

Exposing (*S,E*)-**6** to the cationic gold(I) complex [(JohnPhos)Au(MeCN)]SbF<sub>6</sub> (**13**) for 5 minutes at room temperature gave **7** in 60% yield (Scheme 3). Other gold(I)




**Scheme 3.** Reagents and conditions: a) [(JohnPhos)Au(MeCN)]SbF<sub>6</sub> (**13**; 2 mol%), 23 °C, 5 min (60%); b) H<sub>2</sub>, Pd(OH)<sub>2</sub>/C, 1:1 MeOH/THF, 23 °C, 4 h (79%); c) [Ir(cod)(PCy<sub>3</sub>)py]BAR<sub>F</sub> (15 mol%), H<sub>2</sub> (80 atm), CH<sub>2</sub>Cl<sub>2</sub>, 40 °C, 4 days (40%); d) oxone, NaHCO<sub>3</sub>, 18-crown-6, 1:1:2 acetone/CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O, 23 °C, 1 h (51%); e) Li, EDA, 50 °C, 1 h (78%); f) allyl alcohol (20 equiv), **13** (2 mol%), −30 °C, 15 min (56% + 21% **7**); g) [Pd(PPh<sub>3</sub>)<sub>4</sub>] (5 mol%), K<sub>2</sub>CO<sub>3</sub>, MeOH, reflux 72 h (72%); h) *m*CPBA, CH<sub>2</sub>Cl<sub>2</sub>, 0 to 23 °C (83%); i) Li, EDA, 50 °C, 1.5 h (62%). BAR<sub>F</sub> = 3,5-bis(trifluoromethyl)phenylborate, cod = 1,5-cyclooctadiene, EDA = ethylenediamine, JohnPhos = (2-biphenyl)-di-*tert*-butylphosphine; *m*CPBA = *m*-chloroperbenzoic acid.

catalysts were also screened for this reaction, but the best results were obtained using complex **13**.<sup>[27]</sup> The relative configuration of **7** (racemic series) was confirmed by X-ray diffraction.<sup>[28,29]</sup> Debenzylation of **7** with H<sub>2</sub> (1 atm) and Pd(OH)<sub>2</sub>/C gave the alcohol **14** (79% yield), which was hydrogenated with [Ir(cod)(PCy<sub>3</sub>)py]BAR<sub>F</sub> catalyst<sup>[30]</sup> under high pressure of H<sub>2</sub> to give **3** in 40% yield (95:5 e.r.). The synthesis **3** from **9** required seven steps and proceeded in 12% overall yield.

Epoxidation of **7** with dimethyldioxirane yielded **15** stereoselectively. Epoxide opening and ether cleavage with Li in ethylenediamine<sup>[31]</sup> yielded **5** in 78% (96:4 e.r.), which gave enantiopure material after crystallization. The synthesis of **5** from **9** was accomplished in seven steps with 15% overall yield.

When the gold-catalyzed reaction of dienyne (*S,E*)-**6** was performed in the presence of allyl alcohol as an external nucleophile, the allyl ether **8** was obtained with the opposite configuration at C4 compared to that of **7** (Table 1). While lowering the reaction temperature to −30 °C led to a 1:1 mixture of **7** and **8** (Table 1, entry 3), increasing the concentration of allyl alcohol to 20 equivalents favored the intermolecular pathway (Table 1, entry 5). Similar results were obtained with using only 1 mol% gold(I) catalyst (Table 1, entry 5). Under the optimized reaction conditions, **8** was


**Table 1:** Gold(I)-catalyzed addition of allyl alcohol to (*S,E*)-**6**.<sup>[a]</sup>



Entry	AllylOH (equiv)	T [°C]	t [min]	7/8 <sup>[b]</sup>
1	10	23	5	75:25
2	10	0	10	55:45
3	10	−30	15	50:50
4	20	−30	20	27:73
5 <sup>[c]</sup>	20	−30	30	33:67

[a] 0.05 M. [b] Determined by GC-MS. [c] 1 mol % **13**.

obtained in 56 % yield, along with **7** (21 % yield; Scheme 3). Removal of the allylic ether with [Pd(PPh<sub>3</sub>)<sub>4</sub>] in MeOH gave the alcohol **16**, whose structure was confirmed by X-ray crystal diffraction<sup>[29]</sup> in the racemic series (Figure 2).<sup>[26]</sup> Although **4** could be synthesized from **16**, a more direct



**Figure 2.** X-ray structures for (±)-**16** and **4**. Thermal ellipsoids are shown at 50 % probability.

synthesis was completed from **8** by selective epoxidation with *m*CPBA from the convex face (83 % yield), followed by opening of the epoxide and allyl cleavage with Li in ethylenediamine to give **4** in 62 % yield (87:13 e.r.), yielding enantiopure **4** after crystallization. Spectral data and optical rotation of synthetic 4*α*,7*α*-aromadendranediol matched those reported for the natural compound. The relative and absolute configuration of **4** were confirmed by X-ray diffraction (Figure 2).<sup>[29]</sup> The synthesis of (+)-4*α*,7*α*-aromadendranediol was similarly carried out from (*R,R*)-**10**.<sup>[27]</sup>

The stereochemical divergent synthesis of **3** and **4** from (*S,E*)-**6** confirms the proposal that this cascade cyclization process proceeds by intra- or intermolecular reactions of cyclopropyl gold(I) carbene-like intermediates such as **A** or **B**.<sup>[18,32]</sup> The enantioselectivity is fully preserved in the formation of **3** and **5** via **7** by an intramolecular gold(I)-catalyzed 1,5-migration of a propargylic group. The intermolecular reaction of (*S,E*)-**6** with allyl alcohol occurs with high enantioselectivity (ca. 96 %). In this case, the slight racemization is due to the competitive formation of a propargyl

carbocation, presumably facilitated by the higher polarity of the reaction medium.

In summary, we have completed highly concise syntheses of three representative aromadendranes from a single precursor by a stereodivergent gold-catalyzed reaction which establishes four new stereogenic centers from a single one. The three natural sesquiterpenes (−)-epiglobulol (**3**), (−)-4*α*,7*α*-aromadendranediol (**4**), and (−)-4*β*,7*α*-aromadendranediol (**5**) have been synthesized in seven steps in 12, 17, and 15 % overall yields, respectively, from commercially available (*E,E*)-farnesol (**9**), and constitutes the shortest total syntheses of these natural compounds. This route could be extended for the enantioselective synthesis of any enantiomer of other aromadendranes and non-natural analogues.

Received: February 3, 2014

Published online: April 1, 2014

**Keywords:** cyclization · gold · natural products · terpenoids · total synthesis

- [1] H. J. M. Gijsen, J. B. Wijnberg, A. de Groot, *Prog. Chem. Org. Nat. Prod.* **1995**, *64*, 149–193.
- [2] I. C. Moreira, J. H. G. Lago, N. C. M. Young, N. F. Roque, *J. Braz. Chem. Soc.* **2003**, *14*, 828–831.
- [3] a) C. Gaspar-Marques, M. F. Simões, B. Rodríguez, *J. Nat. Prod.* **2004**, *67*, 614–621; b) Y. Yamakoshi, M. Murata, A. Shimizu, S. Homma, *Biosci. Biotechnol. Biochem.* **1992**, *56*, 1570–1576; c) J. M. Jacyno, N. Montemurro, A. D. Bates, H. G. Cutler, *J. Agric. Food Chem.* **1991**, *39*, 1166–1168; d) M. Murata, Y. Yamakoshi, S. Homma, K. Aida, K. Hori, Y. Ohashi, *Agric. Biol. Chem.* **1990**, *54*, 3221–3226.
- [4] a) M. Nishizawa, M. Emura, Y. Kan, H. Yamada, K. Ogawa, N. Hamanaka, *Tetrahedron Lett.* **1992**, *21*, 2983–2986; b) N. De Tommasi, C. Pizza, C. Conti, N. Orsi, M. L. Stein, *J. Nat. Prod.* **1990**, *53*, 830–835.
- [5] a) Z.-S. Su, S. Yin, Z.-W. Zhou, Y. Wu, J. Diang, J.-M. Yue, *J. Nat. Prod.* **2008**, *71*, 1410–1413; b) H. Tada, F. Yasuda, *Chem. Pharm. Bull.* **1985**, *33*, 1941–1945.
- [6] a) A. Matsuo, K. Atsumi, M. Nakayama, S. Hayashi, *J. Chem. Soc. Perkin Trans. 1* **1981**, 2816–2824; b) T. D. Hubert, D. F. Wiemer, *Phytochemistry* **1985**, *24*, 1197–1198; c) Y. Asakawa, M. Toyota, T. Takemoto, I. Kubo, K. Nakanishi, *Phytochemistry* **1980**, *19*, 2147–2154; d) N. Pérez-Hernández, H. Ponce-Monter, M. I. Ortiz, R. Cariño-Cortés, P. Joseph-Nathan, *Z. Naturforsch. C* **2009**, *64*, 840–846; e) T. S. Wu, Y. Chan, Y. Leu, *Chem. Pharm. Bull.* **2000**, *3*, 357–361.
- [7] a) C. M. Beechan, C. Djerassi, H. Eggert, *Tetrahedron* **1978**, *34*, 2503–2508.
- [8] a) E. Fattorusso, S. Magno, L. Mayol, C. Santacroce, D. Sica, *Tetrahedron Lett.* **1975**, *31*, 269–270; b) C. C. da Silva, V. Almagro, J. Zukerman-Schpector, E. E. Castellano, A. J. Marsaioli, *J. Org. Chem.* **1994**, *59*, 2880–2881; c) S. Kozawa, H. Ishiyama, J. Fromont, J. Kobayashi, *J. Nat. Prod.* **2008**, *71*, 445–447; d) H. Ishiyama, S. Kozawa, K. Aoyama, Y. Mikami, J. Fromont, J. Kobayashi, *J. Nat. Prod.* **2008**, *71*, 1301–1303.
- [9] A. S. R. Anjaneyulu, M. V. R. Krishnamurthy, G. V. Rao, *Tetrahedron* **1997**, *53*, 9301–9312.
- [10] a) G. Büchi, W. Hofheinz, J. V. Paukstelis, *J. Am. Chem. Soc.* **1966**, *88*, 4113–4114; b) G. Buechi, W. Hofheinz, J. V. Paukstelis, *J. Am. Chem. Soc.* **1969**, *91*, 6473–6478; c) J. A. Marshall, J. A. Ruth, *J. Org. Chem.* **1974**, *39*, 1971–1973; d) H. J. M. Gijsen, J. B. P. A. Wijnberg, G. A. Stork, A. de Groot, *Tetrahedron* **1991**,

- 47, 4409–4416; e) H. J. M. Gijsen, J. B. P. A. Wijnberg, G. A. Stork, A. de Groot, M. A. de Waard, J. G. M. van Nistelrooy, *Tetrahedron* **1992**, *48*, 2465–2476; f) H. J. M. Gijsen, J. B. P. A. Wijnberg, C. van Ravenswaay, A. de Groot, *Tetrahedron* **1994**, *50*, 4733–4744; g) H. J. M. Gijsen, J. B. P. A. Wijnberg, A. de Groot, *Tetrahedron* **1994**, *50*, 4745–4754; h) S. L. Gwaltney II, K. J. Shea, *Tetrahedron Lett.* **1996**, *37*, 949–952; i) R. Guillermo, J. R. Hanson, A. Truneh, *J. Chem. Res. Synop.* **1997**, 28–29; j) F. J. Moreno-Dorado, Y. M. A. W. Lamers, G. Mironov, J. B. P. A. Wijnberg, A. de Groot, *Tetrahedron* **2003**, *59*, 7743–7750; k) Y. M. A. W. Lamers, G. Rusu, J. B. P. A. Wijnberg, A. de Groot, *Tetrahedron* **2003**, *59*, 9361–9369.
- [11] R. Tressl, K.-H. Engel, M. Kossa, H. Koppler, *J. Agric. Food Chem.* **1983**, *31*, 892–897.
- [12] a) A. Elaissia, H. Medinia, M. L. Khoubab, M. Simmonds, F. Lynend, F. Farhat, R. Chemli, F. Harzallah-Skhiri, *Chem. Biodiversity* **2011**, *8*, 352–361; b) M. Z. Özel, F. Göğüs, A. C. Lewis, *J. Chromatogr. Sci.* **2008**, *46*, 157–161; c) H. Nazemiyeh, S. M. Razavi, R. Hajiboland, A. Delazar, S. Esna-asharii, R. Bamdad, L. Nahar, S. D. Sarker, *Chem. Nat. Compd.* **2007**, *43*, 736–737; d) M. E. A. Stefanello, A. C. R. F. Pascoal, M. J. Salvador, *Chem. Biodiversity* **2011**, *8*, 73–94.
- [13] H. J. M. Gijsen, K. Kanai, G. A. Stork, J. B. P. A. Wijnberg, R. V. A. Orru, C. G. J. M. Seelen, S. M. van der Kerk, A. de Groot, *Tetrahedron* **1990**, *46*, 7237–7246.
- [14] D. Caine, J. T. Gupton III, *J. Org. Chem.* **1975**, *40*, 809–810.
- [15] Y. Oonishi, A. Taniuchi, M. Mori, Y. Sato, *Tetrahedron Lett.* **2006**, *47*, 5617–5621.
- [16] B. Simmons, A. M. Walji, D. W. C. MacMillan, *Angew. Chem.* **2009**, *121*, 4413–4417; *Angew. Chem. Int. Ed.* **2009**, *48*, 4349–4353.
- [17] Y. Takeda, H. Yamashita, T. Matsumoto, H. Terao, *Phytochemistry* **1993**, *33*, 713–715.
- [18] E. Jiménez-Núñez, M. Raducan, T. Lauterbach, K. Molawi, C. R. Solorio, A. M. Echavarren, *Angew. Chem.* **2009**, *121*, 6268–6271; *Angew. Chem. Int. Ed.* **2009**, *48*, 6152–6155.
- [19] a) E. Jiménez-Núñez, K. Molawi, A. M. Echavarren, *Chem. Commun.* **2009**, 7327–7329; b) K. Molawi, N. Delpont, A. M. Echavarren, *Angew. Chem.* **2010**, *122*, 3595–3597; *Angew. Chem. Int. Ed.* **2010**, *49*, 3517–3519; c) M. Gaydou, R. E. Miller, N. Delpont, J. Ceccon, A. M. Echavarren, *Angew. Chem.* **2013**, *125*, 6524–6527; *Angew. Chem. Int. Ed.* **2013**, *52*, 6396–6399; d) A. Pitaval, D. Leboeuf, J. Ceccon, A. M. Echavarren, *Org. Lett.* **2013**, *15*, 4580–4583.
- [20] A mixture of isomers is also available. (*Z,Z*)- or (*E,Z*)-farnesol are just available as analytical standards. Although (*E,Z*)-farnesol can be prepared in two steps from nerylacetone (10:1 *E,Z/E,E* mixture, 55% yield): J. S. Yu, T. S. Kleckley, D. F. Wiemer, *Org. Lett.* **2005**, *7*, 4803–4806, pure nerylacetone is commercially available only in milligram amounts.
- [21] M. Willot, M. Christmann, *Nat. Chem.* **2010**, *2*, 519–520.
- [22] Selected examples of gold(I)-catalyzed polycyclizations: a) A. Fürstner, L. Morency, *Angew. Chem.* **2008**, *120*, 5108–5111; *Angew. Chem. Int. Ed.* **2008**, *47*, 5030–5033; b) P. Y. Toullec, T. Blarre, V. Michelet, *Org. Lett.* **2009**, *11*, 2888–2891; c) S. G. Sethofer, T. Mayer, F. D. Toste, *J. Am. Chem. Soc.* **2010**, *132*, 8276–8277.
- [23] For lead references on late transition metal catalyzed polycyclizations of polyenes: a) M. A. Schafroth, D. Sarlah, S. Krautwald, E. M. Carreira, *J. Am. Chem. Soc.* **2012**, *134*, 20276–20278; b) O. F. Jeker, A. G. Kravina, E. M. Carreira, *Angew. Chem.* **2013**, *125*, 12388–12391; *Angew. Chem. Int. Ed.* **2013**, *52*, 12166–12169; c) M. J. Geier, M. R. Gagné, *J. Am. Chem. Soc.* **2014**, *136*, 3032–3035.
- [24] D. K. Mohapatra, C. Pramanik, M. S. Chorghade, M. K. Gurjar, *Eur. J. Org. Chem.* **2007**, 5059–5063.
- [25] J. Tanuwidjaja, S.-S. Ng, T. F. Jamison, *J. Am. Chem. Soc.* **2009**, *131*, 12084–12085.
- [26] The syntheses of (±)-epiglobulol and (±)-4α,7α-aromadendranediol was also carried out from a 1.2:1 mixture of geranyl- and nerylacetone. See the Supporting Information for details.
- [27] See the Supporting Information for additional details.
- [28] Racemic **7** was prepared in four steps from a 1.2:1 mixture of geranyl- and nerylacetone.<sup>[27]</sup>
- [29] CCDC 983695 (**4**), 983696 [(±)-**7**], 983697 [(±)-**16**] contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- [30] B. Wüstenberg, A. Pfaltz, *Adv. Synth. Catal.* **2008**, *350*, 174–178.
- [31] H. C. Brown, S. Ikegami, J. H. Kawakami, *J. Org. Chem.* **1970**, *35*, 3243–3245.
- [32] a) E. Jiménez-Núñez, A. M. Echavarren, *Chem. Rev.* **2008**, *108*, 3326–3350; b) C. Obradors, A. M. Echavarren, *Acc. Chem. Res.* **2014**, *47*, 902–912.